

On closed P -sets with ccc in the space ω^*

R.Frankiewicz(Warsaw) S.Shelah(Jerusalem) P.Zbierski(Warsaw)

Abstract. It is proved that – consistently – there can be no ccc closed P -sets in the remainder space ω^* .

In this paper we show how to construct a model of set theory in which there are no P -sets satisfying ccc (countable antichain condition) in the ultrafilter space $\omega^* = \beta\omega \setminus \omega$. The problem of the existence of such sets (which are generalizations of P -points) was known since some time and occurred explicitly in $\mathfrak{c}vM-R$ —. In the proof we follow the construction from $\mathfrak{c}S$ — of a model in which there are no P -points. A particular case of P -sets, which are supports of approximative measures has been settled in $\mathfrak{c}M$ —, where the author shows that there can be no such measures on $P(\omega)/fin$. (Under CH , e.g. the Gleason space $\mathbf{G}(2^\omega)$ of the Cantor space is a ccc P -set in ω^* which carries no approximative measure).

Sec.1. Closed P -sets in the space ω^* can be identified with P -filters F on ω . Thus, the dual ideal $I = \{\omega \setminus A : A \in F\}$ has the property:

$$(1.1) \quad \begin{array}{l} \text{If } A_n \in I, \text{ for } n \in \omega, \text{ then there is an } A \in I \\ \text{such that } A_n \subseteq_* A, \text{ for each } n \in \omega. \end{array}$$

Further, the countable chain condition imposed upon F implies that I is fat in the following sense (see $\mathfrak{c}F-Z$ —):

$$(1.2) \quad \begin{array}{l} \text{if } A_n \in I, \text{ for } n \in \omega \text{ and } \lim_n \min A_n = \infty, \\ \text{then there is an infinite } Z \subseteq \omega \text{ such that } \bigcup_{n \in Z} A_n \in I. \end{array}$$

Indeed, let $e_n = A_n \setminus A$, where $A \in I$ is as in (1.1). Since $\min A_n$ are arbitrarily large, we can find an infinite set $Y \subseteq \omega$ such that the family $\{e_n : n \in Y\}$ is disjoint. If $\{Y_\alpha : \alpha < c\}$ is an almost disjoint family of subsets of Y , then the unions

$$S_\alpha = \bigcup \{e_n : n \in Y_\alpha\}, \quad \alpha < c$$

are almost disjoint and hence the closures S_α^* in the space ω^* are disjoint. By ccc we have

$$S_\alpha^* \cap \bigcap \{B^* : B \in F\} = \emptyset,$$

for some α and consequently $S_\alpha \in I$. It follows that the union

$$\bigcup_{n \in Y_\alpha} A_n = \bigcup_{n \in Y_\alpha} (A_n \cap A) \cup \bigcup_{n \in Y_\alpha} (A_n \setminus A)$$

is in I as a subset of $S_\alpha \cup A$.

Let us fix a given ccc P -filter F and its dual I . We shall define a forcing $\mathbf{P} = \mathbf{P}(F)$.

A partial ordering $(T, \leq_T)$, where $T \subseteq \omega$, will be called a tree, if for each $i \in T$ the set of predecessors $\{j \in T : j \leq_T i\}$ is linearly ordered and

$$i \leq_T j \text{ implies } i \leq j, \text{ for all } i, j \in T.$$

We define a partial ordering for trees

$$\begin{aligned} T \leq_t S &\text{ iff } (S, \leq_S) \text{ is a subordering of } (T, \leq_T) \\ &\text{ and each branch of } T \text{ contains cofinally a (unique) branch of } S. \end{aligned}$$

There is a tree T_0 such that $T_0 \in I$ and T_0 is order isomorphic to the full binary tree of height ω .

Deleting the numbers $\leq n$ from T_0 we obtain a subtree denoted by $T_0^{(n)}$ (we have $T_0^{(n)} \leq_t T_0^{(m)}$, for $n \leq m$). Let $\mathcal{T}$ consist of all the trees $T \in I$ such that

$$T \leq_t T_0^{(n)}, \text{ for some } n \in \omega.$$

Note that each tree $T \in \mathcal{T}$ has finitely many roots.

DEFINITION. *Elements of the forcing $\mathbf{P}$ are of the form $p = \langle T_p, f_p \rangle$, where $T_p \in \mathcal{T}$ and $f_p : T_p \rightarrow \{0, 1\}$. The ordering of $\mathbf{P}$ is defined thus*

$$p \leq q \text{ iff } T_p \leq_t T_q \text{ and } f_p \supseteq f_q.$$

Let $\{b_\alpha : \alpha < c\}$ be a fixed enumeration of all the branches of T_0 in V . For a generic $G \subseteq \mathbf{P}$ let $T_G = \bigcup_{p \in G} T_p$ and $f_G = \bigcup_{p \in G} f_p$.

Each branch B of T_G contains cofinally a unique b_α . Let us write $B = B_\alpha$ and define

$$X_\alpha = \{i \in \omega : i \in B_\alpha \text{ and } f_G(i) = 1\}$$

Since $T_p \in I$, for any $p \in \mathbf{P}$, hence $\omega \setminus T_p \cap A$ is infinite, for each $A \in F$. It follows that the sets

$$D_{n\varepsilon}^{A\alpha} = \{p \in \mathbf{P} : \exists i > n \diamond i \in b_\alpha^p \text{ and } f_p(i) = \varepsilon\}$$

are dense, for each $A \in F$, $n \in \omega$, $\alpha < c$ and $\varepsilon = 0, 1$ (here b_α^p denotes the branch of T_p extending b_α).

Thus, $\mathbf{P}$ adds uncountably many almost disjoint Gregorieff-like sets.

Sec.2. Let $\mathbf{Q} = \mathbf{Q}(F)$ be a countable product of $\mathbf{P} = \mathbf{P}(F)$. Thus the elements $q \in \mathbf{Q}$ can be written in the form

$$q = \langle f_0^q, f_1^q, \dots \rangle, \text{ where } \langle \text{dm}(f_i^q), f_i^q \rangle \in \mathbf{P}, \text{ for each } i < \omega.$$

By $q^{(n)}$ we denote the condition $\langle g_i : i < \omega \rangle$ where

$$g_i = \begin{cases} f_i^q \upharpoonright \text{dm}(f_i^q)^{(n)}, & \text{for } i < n \\ f_i^q & \text{for } i \geq n \end{cases}$$

Here $T^{(n)}$ is a tree obtained from T by deleting the numbers $\leq n$.

LEMMA 2.1. *For each decreasing sequence $p_0 \geq p_1 \geq \dots$ there is a q and an infinite $Z \subseteq \omega$ such that*

$$q \leq p_n^{(n)}, \text{ for each } n \in Z.$$

PROOF. Let $T_{ni} = \text{dm}(f_i^{p_n})$, where $p_n = \langle f_i^{p_n} : i < \omega \rangle$. Since $\min T_{ni}^{(n)} \geq n$, we may use (1.2) to define by induction a descending sequence $Z_0 \supseteq Z_1 \supseteq \dots$ of infinite subsets of ω such that

$$\bigcup_{n \in Z_i} T_{ni}^{(n)} \text{ is in } I, \text{ for each } i < \omega.$$

There is an infinite $Z \subseteq \omega$, such that $Z \subseteq_* Z_i$, for each $i < \omega$. Define

$$T_i = T_{ii} \cup \bigcup_{n \in Z} T_{ni}^{(n)}$$

and

$$f_i^q = f_i^{p_i} \cup \bigcup_{n \in Z} f_i^{p_n} \upharpoonright T_{ni}^{(n)}.$$

Then, $\text{dm}(f_i^q) = T_i$ and $q = \langle f_i^q : i < \omega \rangle$ is as required. *QED*

For $q \in \mathbf{Q}$ and $n \in \omega$ let $S(q, n)$ be the set of all sequences $s = \langle s_0, \dots, s_{n-1} \rangle$ satisfying the following properties

1. $s_0, \dots, s_{n-1}$ are finite zero-one functions.
2. The domains $t_0 = \text{dm}(s_0), \dots, t_{n-1} = \text{dm}(s_{n-1})$ are finite trees such that

$$t_0 \cap T_0^{(n)} = \dots = t_{n-1} \cap T_{n-1}^{(n)} = \emptyset,$$

where $T_0 = \text{dm}(f_0^q), \dots, T_{n-1} = \text{dm}(f_{n-1}^q)$

3. Ordered sums $t_0 \oplus T_0^{(n)}, \dots, t_{n-1} \oplus T_{n-1}^{(n)}$ are trees belonging to $\mathcal{T}$.

Note that from the definition of $\mathcal{T}$ it follows that $S(q, n)$ is always finite. Let us denote

$$s * q^{(n)} = \langle s_0 \cup f_0^q, \dots, s_{n-1} \cup f_{n-1}^q, f_n^q, \dots \rangle$$

for q, n, s as above. Obviously, we have

$$(2.2) \quad \text{the set } \{s * q^{(n)} : s \in S(q, n)\} \text{ is predense below } q^{(n)}$$

(i.e. the boolean sum $\sum_{s \in S(q, n)} s * q^{(n)} = q^{(n)}$).

Now, we obtain easily an analogue of VI, 4.5 in αS —.

$$\begin{aligned}
 & \text{For arbitrary } p \in \mathbf{Q}, n < \omega \text{ and } \tau \in V^{(Q)} \\
 & \text{such that } \mathbf{Q} \Vdash \text{“}\tau \text{ is an ordinal” there is} \\
 (2.3) \quad & \text{a } q \leq p \text{ and ordinals } \{\alpha(s) : s \in S(p, n)\} \\
 & \text{so that} \\
 & q^{(n)} \Vdash \bigvee_s \tau = \alpha(s)
 \end{aligned}$$

Indeed, if $S(q, n) = \{s^0, \dots, s^{m-1}\}$, then we define inductively conditions $p_0, \dots, p_m$ so that $p_0 = p$ and $p_{k+1} \leq s^k * p_k^{(n)}$ is such that

$$p_{k+1} \Vdash \text{“}\tau = \alpha\text{”}, \text{ for some ordinal } \alpha = \alpha(s^k)$$

Now, $q = s * p_m^{(n)}$, where s is such that $p = s * p^{(k)}$ (we may assume $s \in S(p, n)$), satisfies (2.3).

2.4. THEOREM. $\mathbf{Q}$ is α -proper, for every $\alpha < \omega_1$, and has the strong PP-property.

PROOF. Let countable $N \prec H(\kappa)$ for sufficiently large κ , be such that $\mathbf{Q} \in N$ and suppose that $p \in \mathbf{Q} \cap N$. To prove that $\mathbf{Q}$ is proper we have to find a $q \leq p$, which is N -generic. Let $\{\tau_n : n < \omega\}$ be an enumeration of all the $\mathbf{Q}$ -names for ordinals, such that $\tau_n \in N$, for $n < \omega$. Using (2.3) we define inductively a sequence $p_0 = p \geq p_1 \geq \dots$ and ordinals $\alpha(n, s)$ so that

$$p_n^{(n)} \Vdash \bigwedge_{i \leq n} \bigvee_s \tau_i = \alpha(n, s) \text{ for each } n < \omega$$

(i.e. in the n -th step we apply (2.3) for all names $\tau_0, \dots, \tau_n$). Note that the p 's and α 's can be found in N , since $N \prec H(\kappa)$. By Lemma 2.1 there is a q and an infinite $Z \subseteq \omega$ such that

$$q \leq p_n^{(n)}, \text{ for each } n \in Z.$$

Hence also $q^{(m)} \leq p_n^{(n)}$ holds for arbitrarily large n and all $m < \omega$ and thus

$$q^{(m)} \Vdash \text{“}\tau_n \in N\text{”},$$

for all $n, m < \omega$.

By III, 2.6 of αS —, each $q^{(m)}$ is N -generic.

To see that $\mathbf{Q}$ is α -proper let $\langle N_\xi : \xi \leq \alpha \rangle$ be a continuous sequence of elementary countable submodels of $H(\kappa)$ such that $\mathbf{Q} \in N_0$ and

$$\langle N_\xi : \xi \leq \eta \rangle \in N_{\eta+1}, \text{ for each } \eta < \alpha.$$

Assume that $\mathbf{Q}$ is β -proper, for each $\beta < \alpha$ and let $q_0 \in \mathbf{Q} \cap N_0$. If $\alpha = \beta + 1$, we have a $q \leq q_0$ which is N_ξ -generic, for each $\xi \leq \beta$ and we may assume that the $q^{(n)}$ have the same property, for all $n < \omega$. since $N_\alpha \prec H(\kappa)$ and all the parameters are in N_α , such a q can be found in N_α and as above we construct a $q_\alpha \leq q$ which is N_α -generic and so are the $q_\alpha^{(n)}$, for $n < \omega$. Thus, q_α and all the $q_\alpha^{(n)}$ are N_ξ -generic for all $\xi \leq \alpha$. If α is a limit ordinal, we fix an increasing sequence $\langle \xi_n : n < \omega \rangle$ such that $\alpha = \sup_{n < \omega} \xi_n$ and by the inductive hypothesis there is a sequence $q_0 \geq q_{\xi_0} \geq q_{\xi_1} \geq \dots$ such that, for each $n < \omega$, q_{ξ_n} is N_ξ -generic, for each $\xi \leq \xi_n$ and $q_{\xi_n} \in N_{\xi_{n+1}}$ and that $q_{\xi_n}^{(m)}$ have the same property for each $m < \omega$. By Lemma 2.1 there is a $q \in \mathbf{Q}$ such that $q \leq q_{\xi_n}^{(n)}$, for infinitely many $n < \omega$. Thus, $q \leq q_0$ and q is N_ξ -generic for each $\xi < \alpha$ and hence also for each $\xi \leq \alpha$.

Finally, to prove the *PP*-property let $h : \omega \longrightarrow \omega$ diverge to infinity and suppose that $p \Vdash "f : \omega \longrightarrow \omega"$. Define

$$k_n = \min\{i : h(i) > 2^n \cdot \text{card}S(p, n)\}, \text{ for } n < \omega$$

and, using (2.3), define inductively the sequence $p = p_0 \geq p_1 \geq \dots$ such that

$$p_n^{(n)} \Vdash \bigwedge_{i < k_n} \bigvee_{s \in S(p_i, i)} f(i) = \alpha(s, i) \text{ for each } n < \omega \text{ and some integers } \alpha(s, i) < \omega.$$

Let T be the tree built up of integers

$$\{\alpha(s, i) : i < \omega \text{ and } s \in S(p_i, i)\}$$

If $q \leq p_n^{(n)}$, for infinitely many n , then we have $q \Vdash "f \in \text{Lim } T"$ and $T \cap \omega^{k_n}$ has less elements than $h(k_n)$, for all $n < \omega$, which finishes the proof. *QED*

The last point to be discussed is how does $\mathbf{Q} = \mathbf{Q}(F)$ act in the course of iteration.

2.5.LEMMA. *If $\mathbf{R}$ is ω^ω -bounding (i.e. the set of old functions $\omega \longrightarrow \omega$ dominates) and $\mathbf{Q}(F)$ is a complete subforcing of $\mathbf{R}$, then in $V^{(\mathbf{R})}$ the filter F cannot be extended to a ccc P -filter.*

PROOF. Let X_α^n be the α -th set added by n -th factor of the product $\mathbf{Q} = \mathbf{P}^\omega$. Suppose that for some $r \in \mathbf{R}$ and a ccc P -filter $E \in V^{(\mathbf{R})}$ we have

$$r \Vdash "F \subseteq E"$$

Note that for each $n < \omega$, the relation $X_\alpha^n \in E$ hold for at most countably many α 's, since E is ccc. Hence, there is an α such that for all $n < \omega$ we have $\omega \setminus X_\alpha^n \in E$ and, since E is a P -filter, there is an $A \in E$ and a function g , so that $A \subseteq \bigcap_{n < \omega} (\omega \setminus X_\alpha^n) \cup \diamond 0, g(n)$ i.e. for some $r_1 \leq r$ we have

$$(2.6) \quad r_1 \Vdash \bigcap_{n < \omega} (\omega \setminus X_\alpha^n) \cup \diamond 0, g(n) \in E"$$

Since $\mathbf{R}$ is ω^ω -bounding we may assume that $g \in V$. By the assumption, $\mathbf{Q}$ is a complete subforcing of $\mathbf{R}$ and hence there is a $q \in \mathbf{Q}$ such that r is compatible with each $q' \leq q$.

On the other hand, since $T_n = \text{dm}(f_n^q) \in \mathcal{T}$, there is a set $B \in I$ and an increasing sequence $a_0 < a_1 < \dots$ such that $T_n \setminus \diamond 0, a_n) \subseteq B$, $g(n) < a_n$ and $\diamond a_n, a_{n+1}) \setminus B \neq \emptyset$, for each $n < \omega$. Define $q' \leq q$ as follows. For a given n extend T_n by adding elements of $\diamond a_n, a_{n+1}) \setminus B$ on the α -th branch b_α^q and put $f_n^{q'}(i) = 1$, for each $i \in \diamond a_n, a_{n+1}) \setminus B$. Obviously, we have

$$q' \Vdash "(\omega \setminus X_\alpha^n) \cup \diamond 0, g(n)) \cap \diamond a_n, a_{n+1}) \setminus B = \emptyset", \text{ for each } n$$

and hence

$$q' \Vdash " \bigcap_{n < \omega} (\omega \setminus X_\alpha^n) \cup \diamond 0, g(n)) \setminus B \cap \bigcup_{n < \omega} \diamond a_n, a_{n+1}) = \emptyset "$$

Consequently $q' \Vdash " \bigcap_{n < \omega} (\omega \setminus X_\alpha^n) \cup \diamond 0, g(n)) \subseteq_* B "$, which contradicts (2.6). *QED*

The rest of the proof is routine. Beginning with a model V of $2^\omega = \omega_1$ and $2^{\omega_1} = \omega_2$ we iterate with countable supports the forcings $\mathbf{Q}(F)$, for all *ccc* P -filters F booked at each stage $\alpha < \omega_2$ of the iteration. From $\mathfrak{aeS}$ —, V.4 we know that the resulting forcing $\mathbf{R}$ (obtained after ω_2 stages) is proper and ω^ω -bounding. Hence, in $V \diamond G$ there are no *ccc* P -sets.

References

- $\mathfrak{aeF-Z}$ — R.Frankiewicz, P.Zbierski “Strongly discrete subsets in ω^* ” Fund. Math. 129 (1988) pp 173-180
- $\mathfrak{aeM}$ —A.H.Mekler “Finitely additive measures on $\mathbb{N}$ and the additive property” Proc.AMS Vol.92 No3 Nov.1984 pp 439-444
- $\mathfrak{aeV M-R}$ — J.vanMill, G.M.Reed editors “Open problems in topology”, North Holland Elsevier Science Publishers B.V 1990
- $\mathfrak{aeS}$ — S.Shelah “Proper forcing” Lecture Notes in Mathematics 940 Springer Verlag

RYSZARD FRANKIEWICZ Institute of Mathematics, Polish Academy of Sciences, Warsaw, Poland

SAHARON SHELAH Institute of Mathematics, The Hebrew University, Jerusalem, Israel

PAWEŁ ZBIERSKI Institute of Mathematics, Warsaw University, Warsaw, Poland